

Table 1

Summary of the taxonomic changes proposed here

Watson et al. (1986)	Proposed new taxonomy
<i>P. davisoni</i>	
<i>davisoni</i> (Oates, 1889)	<i>P. davisoni</i> monotypic
	<i>P. ogilviegranti</i>
<i>disturbans</i> (La Touche, 1922)	<i>P. o. disturbans</i>
<i>klossi</i> (Riley, 1922)	<i>P. o. klossi</i>
<i>ogilviegranti</i> (La Touche, 1922)	<i>P. o. ogilviegranti</i>
<i>P. reguloides</i>	<i>P. reguloides</i>
<i>reguloides</i> (Blyth, 1842)	<i>P. r. reguloides</i>
<i>kashmiriensis</i> Ticehurst, 1933	<i>P. r. reguloides</i>
<i>assamensis</i> Hartert, 1921	<i>P. r. assamensis</i>
<i>ticehursti</i> Delacour & Greenway, 1939	<i>P. r. ticehursti</i>
<i>claudiae</i> (La Touche, 1922)	<i>P. claudiae</i> monotypic
	<i>P. goodsoni</i>
<i>goodsoni</i> Hartert, 1910	<i>P. g. goodsoni</i>
<i>fokiensis</i> Hartert, 1917	<i>P. g. fokiensis</i>
<i>Seicercus xanthoschistos</i>	<i>Phylloscopus xanthoschistos</i>
<i>xanthoschistos</i> (Gray and Gray, 1846)	<i>P. x. xanthoschistos</i>
<i>tephrodidas</i> Sick, 1939	<i>P. x. tephrodidas</i>

We suggest that *P. davisoni* be treated as two species, *P. davisoni* and *P. ogilviegranti*; *P. reguloides* as three species, *P. reguloides*, *P. claudiae*, and *P. goodsoni*; and that *Seicercus xanthoschistos* be placed in the genus *Phylloscopus*. The taxa *jerdoni* (Brooks, 1871) and *flavogularis* (Godwin-Austen, 1877) are treated as subspecies of *Seicercus xanthoschistos* by Watson et al. (1986), but were not included in the present analysis.