

Phase feeding for growing and finishing pigs

Problem

Large variation between pigs in a litter or production batch makes it difficult to target their nutrient requirements for optimal growth. Diets are therefore formulated with higher lysine and protein contents than recommended, resulting in wastage of nutrients and unnecessarily high N-emissions.

Solution

A phase feeding plan, with two or more phases, will better reflect the actual need of protein and amino acids for pigs at different live weights (growth phases) as dietary content of crude protein and essential amino acids is decreasing with increasing age of the pigs.

Benefits

Phase feeding will more closely match the pig's nutrient requirements and minimise the over- and under-feeding of nutrients. The feed will be better utilised by the pigs, in favour of both production economy and reduced N-emissions.

Practical recommendation

Applicability box

Theme

Pigs, Feeding and ration planning Geographical coverage Global

Application time All year round

Required time Growing/finishing period

Period of impact All year round

Equipment Feed ration planning Best in Growing/finishing period

- To get the maximum benefit from phase feeding, diets and feeding should be established based on actual animal performance and profitability/performance goals for each stage of production. It is easier to develop with a small number of pigs per batch (to manage heterogeneity)
- Diets should be formulated on a digestible amino acid basis rather than on a total amino acid or crude protein basis, crude protein should preferably be kept at a low level and ingredients should be analysed for their nutrient contents.



A single-feed diet meets the nutrient requirements of the pigs "on average" and due to the variation within the group, while 2- or multi-phase feeding will more closely match the pig's nutrient requirements and minimise the over- and under-feeding of nutrients. More phases will better reflect the actual need for protein and amino acids for pigs at different live weights (growth phases). Illustration: Magdalena Presto Åkerfeldt.



PRACTICE ABSTRACT

- A phase feeding system is complex and factors such as the availability of high-quality protein feed ingredients, the managing and ordering of feed as well as the need for additional feed bins on the farm must be considered.
- Consult with an advisor or nutritionist to adjust the feeding plan accordingly to meet the production goals.

Further information

Weblinks

• Check the Organic Farm Knowledge platform for more <u>practical recommendations on pigs</u> as well as <u>feeding</u> and ration planning.

About this practice abstract and OK-Net EcoFeed

Publishers

Department of Animal Nutrition and Management Swedish University of Agricultural Sciences Box 7024, SE-750 07 Uppsala www.slu.se

Research Institute of Organic Agriculture (FiBL) Ackerstrasse 113, Postfach 219, CH-5070 Frick Phone +41 62 865 72 72, info.suisse@fibl.org, www.fibl.org

IFOAM EU Rue du Commerce 124, BE-1000 Brussels Phone +32 2 280 12 23, info@ifoam-eu.org, www.ifoam-eu.org

Phone +52 2 280 12 23, into@itoani-eu.org, www.itoani-eu.o

Author: Magdalena Presto Åkerfeldt (SLU) Contact: magdalena.akerfeldt@slu.se

Permalink: Organic-farmknowledge.org/tool/35451



OK-Net EcoFeed: This practice abstract was elaborated in the Organic Knowledge Network on Monogastric Animal Feed project. The project is running from January 2018 to December 2020. The overall aim of OK-Net EcoFeed is to help farmers, breeders and the organic feed processing industry in achieving the goal of 100% use of organic and regional feed for monogastrics.

Project website: ok-net-ecofeed.eu

Project partners: IFOAM EU Group (project coordinator), BE; Aarhus University (ICROFS), DK; Organic Research Centre (ORC), UK; Institut Technique de l'Agriculture Biologique (ITAB), FR; Research Institute of Organic Agriculture (FiBL), CH; Bioland, DE; Associazione Italiana per l'Agricoltura Biologica (AIAB), IT; Donau Soja DS, AT; Swedish University of Agricultural Sciences, SE; ECOVALIA, ES; Soil Association, UK. © 2020

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 773911. This communication only reflects the author's view. The Research Executive Agency is not responsible for any use that may be made of the information provided. The authors and editors do not assume responsibility or liability for any possible factual inaccuracies or damage resulting from the application of the recommendations in this practice abstract.

